

NUCLEOTIDE SEQUENCES CODING FOR POLYPEPTIDES ENDOWED WITH A
LARVICIDAL ACTIVITY TOWARDS LEPIDOPTERA

5 The subject of the invention is nucleotide sequences coding
for polypeptides endowed with a larvicidal activity towards
Lepidoptera.

10 It relates more particularly to agents, in particular
nucleotide sequences, polypeptides or even vectors, or bacterial
strains modified by these sequences and expressing polypeptides making
it possible to prepare larvicidal compositions active against
Lepidoptera, preferably against Spodoptera littoralis (hereafter
S.littoralis) or Mamestra brassicae (hereafter designated by
M.brassicae) or capable of transforming the plants to be treated in
15 conferring on them this type of activity.

 It is known that most of the isolates of B.thuringiensis
show a toxic activity with regard to larvae of more than a hundred
species of Lepidoptera.

20 This activity results from the capacity of the strains of
B.thuringiensis to synthesize, at the moment of sporulation,
crystalline inclusions of protein nature, or δ -endotoxins, under the
control of one or several types of gene.

 It has been shown that the activity of these polypeptides
is contained in the NH_2 -terminal half or N-terminus of the protein.

25 The studies carried out have shown the high specificity
of the δ -endotoxins towards larvae of a given species.

 On account of this high specificity, many species of
Lepidoptera, in particular of the family of the Noctuidae, react only
weakly to commercial preparations of available B.thuringiensis.

30 It is so in particular for the species S.littoralis, a poly-
phagous insect which constitutes the principal parasite of cotton
and other industrially important crops. Among these crops, mention
should be made of maize, the castor oil plant, tobacco, the groundnut,
fodder plants, such as clover or alfalfa, or also market garden produce
35 such as the cabbage or the tomato.

filed
12/11/89
6/5/95

Hence, one can imagine the interest of disposing of agents targeting specifically and effectively the family of the Noctuidae and in particular S.littoralis or M.brassicae.

The genes for δ -endotoxins hitherto identified do not code for a polypeptide preferentially active with regard to S.littoralis.

The search by the inventors for a sequence of nucleotides coding for a polypeptide preferably active against the Noctuidae, more especially against S.littoralis, has led them to study the natural isolates of two strains of B.thuringiensis, the larvicidal activity of which on S.littoralis appears to be higher than that of the industrial preparations made starting from other strains of B.thuringiensis.

The species in question are aizawai 7-29 and entomocidus 6-01.

The study of these isolates has made it possible to demonstrate the existence of several genes for δ -endotoxins of different structures and different specificities, of which two genes preferentially active against P.brassicae but not very active against the Noctuida of cotton and a gene inactive against P.brassicae and S.littoralis.

By studying the total DNA of these isolates and by carrying out appropriate hybridizations, followed by the cloning of the fragments identified by hybridization, the inventors have observed that it is possible to isolate nucleotide sequences implicated in genes for δ -endotoxins coding for polypeptides active, preferably, against S.littoralis.

Thus, the aim of the invention is to provide nucleotide sequences capable of coding for at least the NH₂-terminal part of a δ -endotoxin toxic against the Noctuidae and preferably against S.littoralis or M.brassicae.

It also has the aim of providing a polypeptide toxic with regard to the Noctuidae.

Furthermore, the invention relates to a procedure for obtaining such a sequence and a polypeptide showing the desired activity as well as the intermediate agents such as vectors and

00919495.000101

bacterial strains which can be utilized for obtaining the polypeptide.

In addition, the invention relates to the uses of these sequences and polypeptides for the development of larvicidal compositions with regard to the Noctuidae, in particular S.littoralis and for the transformation of the plants likely to be infected by these larvae.

The invention relates to a sequence of nucleotides coding for at least a part of the N-terminal region of a polypeptide toxic specifically against the larvae of Lepidoptera of the Noctuidae family, and preferably against S.littoralis, characterized by its capacity of hybridization with a gene capable of expressing a polypeptide toxic towards larvae of S.littoralis.

According to another aspect of the invention, the nucleotide sequence is characterized in that it is carried by a sequence of nucleotides of about 3 kb such as obtained by in vitro genetic recombination of sequences of nucleotides of B.thuringiensis capable of hybridizing with probes 1, 2 and 3 of pHTA2 shown in figure 2. The fragment of 3 kb corresponds more particularly to the restriction fragment HindIII-PstI.

The sequences of nucleotides of the invention are, in addition, characterized in that they contain sites in the following order : HindIII - HincII - BglIII - KpnI - HindIII - PstI.

In a preferred manner, these sequences of nucleotides are obtained by in vitro genetic recombination of DNA sequences derived from at least one strain of B.thuringiensis. In a variant of the embodiment of the invention, two different strains of B.thuringiensis are utilized.

Strains of B.thuringiensis particularly suited for obtaining these sequences of nucleotides are the strains corresponding to aizawai 7-29 and entomocidus 6-01, deposited on 21 April 1987 under the No. I-661 and No. I-660, respectively, with the National Collection of Cultures of Microorganisms (N.C.C.M.) in Paris.

In an advantageous manner, the sequences of nucleotides of the invention code for a polypeptide capable of forming an immunological complex with antibodies directed against polypeptides showing the larvicidal activity with regard to S.littoralis.

INS
AI

4

A sequence of nucleotides according to the invention is characterized in that it has the capacity to hybridize with a probe formed from the sequence (I) showing the following chain arrangement:

GTC ⁵²TAC TTG ACA GCG GTA GCA ACA TAA TCG GTC AAT TTT AAA
TAT GGG GCA TAT ATT GAT ATT ¹¹²TTA TAA AAT TTG TTA CGT TTT
TTG TAT TTT TTC ATA AGA TGT GTC ATA TGT ATT AAA TCG ¹⁷²TGG
TAA TGA AAA ACA GTA ²³²TCA AAC TAT CAG AAC TTT GGT AGT TTA
ATA AAA AAA CCG AGG TAT TTT ATG GAG GAA AAT ²⁹²AAT CAA AAT
CAA TGC ATA CCT TAC AAT TGT TTA AGT AAT CCT ²⁹²GAA GAA GTA
CTT TTG GAT ³⁵²GGA GAA CCG ATA TCA ACT GGT AAT TCA TCA ATT
GAT ATT TCT CTG TCA CTT GTT CAG TTT CTG ⁴¹²GTA TCT AAC TTT
GTA CCA GGG GGA GGA TTT TTA GTT GCA ⁴¹²TTA ATA GAT TTT GTA
TGG ⁴⁷²GGA ATA GTT GGC CCT TCT CAA TGG GAT GCA TTT CTA GTA
CAA ATT GAA CAA TTA ATT AAT ⁵³²GAA AGA ATA GCT GAA TTT GCT
AGG AAT GCT GCT ATT GCT AAT ⁵³²TTA GAA GGA TTA GGA AAC AAT
TTC AAT ATA TAT GTG GAA GCA TTT AAA GAA TGG GAA ⁵⁹²GAA GAT
CCT AAT AAT CCA GAA ACC AGG ACC AGA ⁶⁵²GTA ATT GAT CGC TTT
CGT ATA CTT GAT GGG ⁶⁵²CTA CTT GAA AGG GAC ATT ⁷¹²CCT TCG TTT
CGA ATT TCT GGA TTT GAA GTA CCC CTT TTA TCC ⁷¹²GTT TAT GCT
CAA GCG GCC ⁷⁷²AAT CTG CAT CTA GCT ATA TTA AGA GAT TCT GTA
ATT TTT GGA GAA AGA TGG GGA TTG ACA ACG ATA AAT GTC AAT
GAA AAC TAT AAT AGA CTA ATT AGG CAT ⁸³²ATT GAT GAA TAT GCT
GAT ⁸⁹²CAC TGT GCA AAT ACG TAT AAT CGG GGA TTA AAT AAT TTA
CCG ⁸⁹²AAA TCT ACG TAT CAA GAT TGG ATA ACA TAT AAT CGA TTA
CGG AGA GAC TTA ACA TTG ⁹⁵²ACT GTA TTA GAT ATC GCC GCT TTC
TTT CCA AAC TAT GAC

00940405.000404

Sequences of nucleotides coding for at least a part of the N-terminal region of a polypeptide toxic specifically towards larvae of Lepidoptera of the Noctuidae family, and preferably towards S.littoralis, are characterized in that they contain the chain arrangement (I) defined above.

In an advantageous manner, the sequence of nucleotides characterized by the chain arrangement defined above codes for a part of a polypeptide having a higher larvicidal activity towards S.littoralis than that of the polypeptides encoded by natural isolates presently known for their effects against S.littoralis.

The study of this sequence of nucleotides shows that it is characterized by an initiation codon ATG situated at position 241 starting from which an open reading frame of 750 nucleotides has been identified.

This sequence is also characterized by a GGAGG attachment site for ribosomes at positions 230 to 234.

According to another feature, the sequence of nucleotides of the invention is characterized in that it contains, upstream from the ATG codon, a sequence going from the nucleotide at position 137 to the nucleotide at position 177, strongly homologous with the region found by Wong et al. (1983) and described in (16) upstream from the gene for the crystal of the strain kurstaki HD1 Dipel (BTK) and for which the authors have shown that it contains three promoters BtI, BtII and Ec which are functional in B.thuringiensis and E.coli, respectively. The homology of these sequences is about 70%.

The invention also relates to a sequence of nucleotides coding for the following sequence (II) of amino acids :

25
TWS
A
30

MET GLU GLU ASN ASN GLN ASN
GLN CYS ILE PRO TYR ASN CYS LEU SER ASN PRO GLU GLU VAL
LEU LEU ASP GLY GLU ARG ILE SER THR GLY ASN SER SER ILE
ASP ILE SER LEU SER LEU VAL GLN PHE LEU VAL SER ASN PHE
VAL PRO GLY GLY PHE LEU VAL GLY LEU ILE ASP PHE VAL TRP
GLY ILE VAL GLY PRO SER GLN TRP ASP ALA PHE LEU VAL GLN
ILE GLU GLN LEU ILE ASN GLU ARG ILE ALA GLU PHE ALA ARG
ASN ALA ALA ILE ALA ASN LEU GLU GLY LEU GLY ASN ASN PHE
ASN ILE TYR VAL GLU ALA PHE LYS GLU TRP GLU GLU ASP PRO
ASN ASN PRO GLU THR ARG THR ARG VAL ILE ASP PRG PHE ARG
ILE LEU ASP GLY LEU LEU GLU ARG ASP ILE PRO SER PHE ARG
ILE SER GLY PHE GLU VAL PRO LEU LEU SER VAL TYR ALA GLN
ALA ALA ASN LEU HIS LEU ALA ILE LEU ARG ASP SER VAL ILE
PHE GLY GLU ARG TRP GLY LEU THR THR ILE ASN VAL ASN GLU
ASN TYR ASN ARG LEU ILE ARG HIS ILE ASP GLU TYR ALA ASP
HIS CYS ALA ASN THR TYR ASN ARG GLY LEU ASN ASN LEU PRO
LYS SER THR TYR GLN ASP TRP ILE THR TYR ASN ARG LEU ARG
ARG ASP LEU THR LEU THR VAL LEU ASP ILE ALA ALA PHE PHE
PRO ASN TYR ASP

F010005 504000

A better identification of the sequence of nucleotides isolated from the above strains, deposited with the NCCM has made it possible to observe that the nucleotide situated at position 273 is guanine (G), the amino acid resulting from the GTA codon thus being valine.

5

Now, the reading of the nucleotide corresponding to this position 273 in the application FR.8708090 of 10 June 1987 had led to reporting thymine (T) and leucine as amino acid resulting from the TTA codon.

Another sequence of nucleotides of the invention is characterized by its capacity of hybridization with a probe formed from the sequence (III) showing the following chain arrangement:

INS¹⁰
A³

15

20

25

30

35

FR 88 09812

FOURTH SET

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

FEDERAL BUREAU OF INVESTIGATION

[illegible]

1641
 1901
 2071
 2161
 2231
 2301
 2371
 2441
 2511
 2581
 2651
 2721
 2791
 2861
 2931
 3001
 3071
 3141
 3211
 3281
 3351
 3421
 3491
 3561
 3631
 3701
 3771
 3841
 3911
 3981
 4051
 4121
 4191
 4261
 4331
 4401
 4471
 4541
 4611
 4681
 4751
 4821
 4891
 4961
 5031
 5101
 5171
 5241
 5311
 5381
 5451
 5521
 5591
 5661
 5731
 5801
 5871
 5941
 6011
 6081
 6151
 6221
 6291
 6361
 6431
 6501
 6571
 6641
 6711
 6781
 6851
 6921
 6991
 7061
 7131
 7201
 7271
 7341
 7411
 7481
 7551
 7621
 7691
 7761
 7831
 7901
 7971
 8041
 8111
 8181
 8251
 8321
 8391
 8461
 8531
 8601
 8671
 8741
 8811
 8881
 8951
 9021
 9091
 9161
 9231
 9301
 9371
 9441
 9511
 9581
 9651
 9721
 9791
 9861
 9931
 10001
 10071
 10141
 10211
 10281
 10351
 10421
 10491
 10561
 10631
 10701
 10771
 10841
 10911
 10981
 11051
 11121
 11191
 11261
 11331
 11401
 11471
 11541
 11611
 11681
 11751
 11821
 11891
 11961
 12031
 12101
 12171
 12241
 12311
 12381
 12451
 12521
 12591
 12661
 12731
 12801
 12871
 12941
 13011
 13081
 13151
 13221
 13291
 13361
 13431
 13501
 13571
 13641
 13711
 13781
 13851
 13921
 13991
 14061
 14131
 14201
 14271
 14341
 14411
 14481
 14551
 14621
 14691
 14761
 14831
 14901
 14971
 15041
 15111
 15181
 15251
 15321
 15391
 15461
 15531
 15601
 15671
 15741
 15811
 15881
 15951
 16021
 16091
 16161
 16231
 16301
 16371
 16441
 16511
 16581
 16651
 16721
 16791
 16861
 16931
 17001
 17071
 17141
 17211
 17281
 17351
 17421
 17491
 17561
 17631
 17701
 17771
 17841
 17911
 17981
 18051
 18121
 18191
 18261
 18331
 18401
 18471
 18541
 18611
 18681
 18751
 18821
 18891
 18961
 19031
 19101
 19171
 19241
 19311
 19381
 19451
 19521
 19591
 19661
 19731
 19801
 19871
 19941
 20011
 20081
 20151
 20221
 20291
 20361
 20431
 20501
 20571
 20641
 20711
 20781
 20851
 20921
 20991
 21061
 21131
 21201
 21271
 21341
 21411
 21481
 21551
 21621
 21691
 21761
 21831
 21901
 21971
 22041
 22111
 22181
 22251
 22321
 22391
 22461
 22531
 22601
 22671
 22741
 22811
 22881
 22951
 23021
 23091
 23161
 23231
 23301
 23371
 23441
 23511
 23581
 23651
 23721
 23791
 23861
 23931
 24001
 24071
 24141
 24211
 24281
 24351
 24421
 24491
 24561
 24631
 24701
 24771
 24841
 24911
 24981
 25051
 25121
 25191
 25261
 25331
 25401
 25471
 25541
 25611
 25681
 25751
 25821
 25891
 25961
 26031
 26101
 26171
 26241
 26311
 26381
 26451
 26521
 26591
 26661
 26731
 26801
 26871
 26941
 27011
 27081
 27151
 27221
 27291
 27361
 27431
 27501
 27571
 27641
 27711
 27781
 27851
 27921
 27991
 28061
 28131
 28201
 28271
 28341
 28411
 28481
 28551
 28621
 28691
 28761
 28831
 28901
 28971
 29041
 29111
 29181
 29251
 29321
 29391
 29461
 29531
 29601
 29671
 29741
 29811
 29881
 29951
 30021
 30091
 30161
 30231
 30301
 30371
 30441
 30511
 30581
 30651
 30721
 30791
 30861
 30931
 31001
 31071
 31141
 31211
 31281
 31351
 31421
 31491
 31561
 31631
 31701
 31771
 31841
 31911
 31981
 32051
 32121
 32191
 32261
 32331
 32401
 32471
 32541
 32611
 32681
 32751
 32821
 32891
 32961
 33031
 33101
 33171
 33241
 33311
 33381
 33451
 33521
 33591
 33661
 33731
 33801
 33871
 33941
 34011
 34081
 34151
 34221
 34291
 34361
 34431
 34501
 34571
 34641
 34711
 34781
 34851
 34921
 34991
 35061
 35131
 35201
 35271
 35341
 35411
 35481
 35551
 35621
 35691
 35761
 35831
 35901
 35971
 36041
 36111
 36181
 36251
 36321
 36391
 36461
 36531
 36601
 36671
 36741
 36811
 36881
 36951
 37021
 37091
 37161
 37231
 37301
 37371
 37441
 37511
 37581
 37651
 37721
 37791
 37861
 37931
 38001
 38071
 38141
 38211
 38281
 38351
 38421
 38491
 38561
 38631
 38701
 38771
 38841
 38911
 38981
 39051
 39121
 39191
 39261
 39331
 39401
 39471
 39541
 39611
 39681
 39751
 39821
 39891
 39961
 40031
 40101
 40171
 40241
 40311
 40381
 40451
 40521
 40591
 40661
 40731
 40801
 40871
 40941
 41011
 41081
 41151
 41221
 41291
 41361
 41431
 41501
 41571
 41641
 41711
 41781
 41851
 41921
 41991
 42061
 42131
 42201
 42271
 42341
 42411
 42481
 42551
 42621
 42691
 42761
 42831
 42901
 42971
 43041
 43111
 43181
 43251
 43321
 43391
 43461
 43531
 43601
 43671
 43741
 43811
 43881
 43951
 44021
 44091
 44161
 44231
 44301
 44371
 44441
 44511
 44581
 44651
 44721
 44791
 44861
 44931
 45001
 45071
 45141
 45211
 45281
 45351
 45421
 45491
 45561
 45631
 45701
 45771
 45841
 45911
 45981
 46051
 46121
 46191
 46261
 46331
 46401
 46471
 46541
 46611
 46681
 46751
 46821
 46891
 46961
 47031
 47101
 47171
 47241
 47311
 47381
 47451
 47521
 47591
 47661
 47731
 47801
 47871
 47941
 48011
 48081
 48151
 48221
 48291
 48361
 48431
 48501
 48571
 48641
 48711
 48781
 48851
 48921
 48991
 49061
 49131
 49201
 49271
 49341
 49411
 49481
 49551
 49621
 49691
 49761
 49831
 49901
 49971
 50041
 50111
 50181
 50251
 50321
 50391
 50461
 50531
 50601
 50671
 50741
 50811
 50881
 50951
 51021
 51091
 51161
 51231
 51301
 51371
 51441
 51511
 51581
 51651
 51721
 51791
 51861
 51931
 52001
 52071
 52141
 52211
 52281
 52351
 52421
 52491
 52561
 52631
 52701
 52771
 52841
 52911
 52981
 53051
 53121
 53191
 53261
 53331
 53401
 53471
 53541
 53611
 53681
 53751
 53821
 53891
 53961
 54031
 54101
 54171
 54241
 54311
 54381
 54451
 54521
 54591
 54661
 54731
 54801
 54871
 54941
 55011
 55081
 55151
 55221
 55291
 55361
 55431
 55501
 55571
 55641
 55711
 55781
 55851
 55921
 55991
 56061
 56131
 56201
 56271
 56341
 56411
 56481
 56551
 56621
 56691
 56761
 56831
 56901
 56971
 57041
 57111
 57181
 57251
 57321
 57391
 57461
 57531
 57601
 57671
 57741
 57811
 57881
 57951
 58021
 58091
 58161
 58231
 58301
 58371
 58441
 58511
 58581
 58651
 58721
 58791
 58861
 58931
 59001
 59071
 59141
 59211
 59281
 59351
 59421
 59491
 59561
 59631
 59701
 59771
 59841
 59911
 59981
 60051
 60121
 60191
 60261
 60331
 60401
 60471
 60541
 60611
 60681
 60751
 60821
 60891
 60961
 61031
 61101
 61171
 61241
 61311
 61381
 61451
 61521
 61591
 61661
 61731
 61801
 61871
 61941
 62011
 62081
 62151
 62221
 62291
 62361
 62431
 62501
 62571
 62641
 62711
 62781
 62851
 62921
 62991
 63061
 63131
 63201
 63271
 63341
 63411
 63481
 63551
 63621
 63691
 63761
 63831
 63901
 63971
 64041
 64111
 64181
 64251
 64321
 64391
 64461
 64531
 64601
 64671
 64741
 64811
 64881
 64951
 65021
 65091
 65161
 65231
 65301
 65371
 65441
 65511
 65581
 65651
 65721
 65791
 65861
 65931
 66001
 66071
 66141
 66211
 66281
 66351
 66421
 66491
 66561
 66631
 66701
 66771
 66841
 66911
 66981
 67051
 67121
 67191
 67261
 67331
 67401
 67471
 67541
 67611
 67681
 67751
 67821
 67891
 67961
 68031
 68101
 68171
 68241
 68311
 68381
 68451
 68521
 68591
 68661
 68731
 68801
 68871
 68941
 69011
 69081
 69151
 69221
 69291
 69361
 69431
 69501
 69571
 69641
 69711
 69781
 69851
 69921
 70001
 70071
 70141
 70211
 70281
 70351
 70421
 70491
 70561
 70631
 70701
 70771
 70841
 70911
 70981
 71051
 71121
 71191
 71261
 71331
 71401
 71471
 71541
 71611
 71681
 71751
 71821
 71891
 71961
 72031
 72101
 72171
 72241
 72311
 72381
 72451
 72521
 72591
 72661
 72731
 72801
 72871
 72941
 73011
 73081
 73151
 73221
 73291
 73361
 73431
 73501
 73571
 73641
 73711
 73781
 73851
 73921
 73991
 74061
 74131
 74201
 74271
 74341
 74411
 74481
 74551
 74621
 74691
 74761
 74831
 74901
 74971
 75041
 75111
 75181
 75251
 75321
 75391
 75461
 75531
 75601
 75671
 75741
 75811
 75881
 75951
 76021
 76091
 76161
 76231
 76301
 76371
 76441
 76511
 76581
 76651
 76721
 76791
 76861
 76931
 77001
 77071
 77141
 77211
 77281
 77351
 77421
 77491
 77561
 77631
 77701
 77771
 77841
 77911
 77981
 78051
 78121
 78191
 78261
 78331
 78401
 78471
 78541
 78611
 78681
 78751
 78821
 78891
 78961
 79031
 79101
 79171
 79241
 79311
 79381
 79451
 79521
 79591
 79661
 79731
 79801
 79871
 79941
 80011
 80081
 80151
 80221
 80291
 80361
 80431
 80501
 80571
 80641
 80711
 80781
 80851
 80921
 80991
 81061
 81131
 81201
 81271
 81341
 81411
 81481
 81551
 81621
 81691
 81761
 81831
 81901
 81971
 82041
 82111
 82181
 82251
 82321
 82391
 82461
 82531
 82601
 82671
 82741
 82811
 82881
 82951
 83021
 83091
 83161
 83231
 83301
 83371
 83441
 83511
 83581
 83651
 83721
 83791
 83861
 83931
 84001
 84071
 84141
 84211
 84281
 84351
 84421
 84491
 84561
 84631
 84701
 84771
 84841
 84911
 84981
 85051
 85121
 85191
 85261
 85331
 85401
 85471
 85541
 85611
 85681
 85751
 85821
 85891
 85961
 86031
 86101
 86171
 86241
 86311
 86381
 86451
 86521
 86591
 86661
 86731
 86801
 86871
 86941
 87011
 87081
 87151
 87221
 87291
 87361
 87431
 87501
 87571
 87641
 87711
 87781
 87851
 87921

5

10

15

20

25

30

INS 4
A

The invention also relates to a sequence of nucleotides characterized in that it codes for a polypeptide containing the sequence (IV) of amino acids below:

W O S S / O

12

271
PRO TYR ASN CYS LEU SER ASN PRO GLU VAL LEU LEU ASP GLY GLU ARG ILE SER THR GLY ASN SER SER ILE ASP ILE SER LEU SER
361
LEU VAL GLN PHE LEU VAL SER ASN PHE VAL PRO GLY GLY PHE LEU VAL GLY LEU ILE ASP PHE VAL IAP GLY ILE VAL GLY PRO SER
431
GLN IAP ASP ALA PHE LEU VAL GLN ILE GLU GLN LEU ILE ASN GLU ARG ILE ALA GLU PHE ALA ARG ASN ALA ALA ILE ALA ASN LEU GLU
541
GLY LEU GLY ASN ASN PHE ASN ILE TYR VAL GLU ALA PHE LYS GLU IAP GLU ASP PRO ASN ASN PRO ALA THR ARG THR ARG VAL ILE
631
ASP ARG PHE ARG ILE LEU ASP GLY LEU LEU GLU ARG ASP ILE PRO SER PHE ARG ILE SER GLY PHE GLU VAL PRO LEU LEU SER VAL TYR
721
ALA GLN ALA ALA ASN LEU HIS LEU ALA ILE LEU ARG ASP SER VAL ILE PHE GLY GLU ARG IAP GLY LEU THR IMA ILE ASN VAL ASN GLU
811
ASA TYR ASN ARG LEU ILE ARG HIS ILE ASP GLU TYR ALA ASP HIS CYS ALA ASN THR TYR ASN ARG GLY LEU ALN ASN LEU PRO LYS SER
901

991

ASH ARG ARG TYR PRO ILE GLN PRO VAL GLY GLN LEU THR ARG GLU VAL TYR THR ASP PRO LEU ILE ASH PHE ASH PRO GLN LEU GLN SER
 1001 VAL ALA GLN LEU PRO THR PHE ASH VAL MET GLU SER SER ALA ILE ARG ASH PRO HIS LEU PHE ASP ILE LEU ASH ASH LEU THR ILE PHE
 1171 THR ASP TRP PHE SER VAL GLY ARG ASH PHE TYR IAP GLY GLY HIS ARG VAL ILE SER SER LEU ILE GLY GLY ASH ILE THR SER PRO
 1261 ILE TYR GLY ARG GLU ALA ASH GLN GLU PRO PRO ARG SER PHE THR PHE ASH GLY PRO VAL PHE ARG THR LEU SER ILE PRO THR LEU ARG
 1331 LEU LEU GLN PRO CYS GLN ARG HIS PHE ASH LEU ARG GLY GLY GLU VAL GLU PHE SER THR PRO THR ASH SER PHE THR ILE
 1441 ARG GLY ARG GLY THR VAL ASP SER LEU THR GLU LEU PRO PRO GLU ASP ASH SER VAL PRO PRO ARG GLU TYR SER HIS ARG LEU CYS
 1531 HIS ALA THR PHE VAL GLN ARG SER GLY THR PRO PHE LEU THR THR GLY VAL VAL PHE SER TRP THR HIS ARG SER ALA THR LEU THR ASH
 1621 THR ILE ASP PRO GLU ARG ILE ASH GLN ILE PRO LEU VAL LYS GLY PHE ARG VAL TRP GLY THR SER VAL ILE THR GLY PRO GLY PHE
 1711 THR GLY GLY ASP ILE LEU ARG ARG ASH THR PHE GLY ASP PHE VAL SER LEU GLN VAL ASH ILE ASH SER PRO ILE THR GLN ARG TYR ARG
 1801 LEU ARG PHE ARG TYR ALA SER SER ARG ASP ALA ARG VAL ILE VAL LEU THR GLY ALA ALA SER THR GLY VAL GLY GLN VAL SER VAL

00010105 50101000

1111 PRO LEU GLN LYS THR MET GLU ILE GLY, GLU ASN LEU THR SER ARG THR PHE ARG TYR THR ASP PHE SER, ASN PRO PHE SER PHE
1201 ARG ALA ASN PRO ASP ILE ILE GLY ILE SER GLU GLN PRO LEU PHE GLY ALA GLY SER ILE SER SER GLY GLU LEU TYR ILE ASP LYS ILE
2071 GLU ILE ILE LEU ALA ASP ALA THR PHE GLU ALA, GLU SER ASP LEU GLU ARG ALA GLN LYS ALA VAL ASN ALA LEU PHE THR SER SER ASN
2101 GLN ILE GLY LEU LYS THR ASP VAL THR ASP TYR, HIS ILE ASP GLN VAL SER ASN LEU VAL ASP CYS LEU SER ASP GLU PHE CYS LEU ASP
2231 GLU LYS ARG GLU LEU SER GLU LYS VAL LYS HIS ALA LYS ARG LEU SER ASP, GLU ARG ASN LEU LEU GLN ASP PRO ASN PHE ARG GLY ILE
2341 ASN ARG GLN PRO ASP ARG GLY TRP ARG GLY SER THR ASP ILE THR ILE GLN GLY GLY ASP VAL PHE LYS GLU ASN TYR VAL THR LEU
2431 PRO GLY THR VAL ASP GLU CYS TYR PRO THR TYR LEU TYR GLN LYS ILE ASP GLU SER LYS LEU LYS ALA TYR THR ARG TYR GLU LEU ARG
2521 GLY TYR ILE GLU ASP SER GLN ASP LEU GLU ILE TYR LEU ILE ALA TYR ASN ALA LYS HIS GLU ILE VAL ASN VAL PRO GLY THR GLY SER
2611 LEU TRP PRO LEU SER ALA GLN SER PRO ILE GLY LYS CYS GLY GLU PRO ASN ARG CYS ALA PRO HIS LEU GLU TRP ASN PRO ASP LEU ASP
2701 CYS SER CYS

The invention also relates to recombinant expression and cloning vectors comprising more particularly at least one sequence of nucleotides such as that defined above, in particular at least a part of the sequence of about 3 kb.

5 A specific recombinant vector is, for example, a plasmid containing the HindIII-PstI fragment of the sequence of nucleotides of the invention, inserted in a vector pUC9. A first preferred vector is the plasmid pHT71, the construction of which is reported in the assemblies below, which comprises a HindIII-PstI DNA fragment according
10 to the invention constituted uniquely of DNA derived from the strain aizawai 7-29.

Another recombinant vector is constituted by the plasmid pHT 671, the construction of which is given in figure 4. This plasmid contains a chimeric HindIII-PstI fragment, obtained by fusing a HindIII-HindII fragment of 1.1 kb derived from the strain entomocidus 6-01
15 with a HincII-PstI fragment of 1.9 kb derived from the strain aizawai 7-29.

The modified bacterial strains which contain one of the nucleotide sequences defined above or also a recombinant expression vector and cloning previously defined, and preferably the plasmid pHT671 or the plasmid pHT71, also enter into the framework of the
20 invention.

The invention also relates to a polypeptide toxic towards larvae of Lepidoptera and in a preferential manner towards
25 S.littoralis, which attack cotton leaves or other crops such as those listed above, characterized in that it is capable of forming an immunological complex with antibodies directed against polypeptides with larvicidal activity towards S.littoralis.

The invention relates more particularly to the NH₂-terminal part of this polypeptide which contains the larvicidal activity.
30

The extremity of the active NH₂-terminal part corresponds to the sequence (II) of amino acids given above.

A preferred polypeptide of the invention is that which corresponds to this sequence (II) and corresponds to the sequence
35 (IV) of amino acids given in the preceding pages. This polypeptide corresponding to the sequence (IV) contains 823 amino acids. Its calculated molecular mass is 92906 Da.

09812-00292-15

This sequence of δ -endotoxin was compared with amino acid sequences of δ -endotoxins derived from other strains of B.thuringiensis active on the Lepidoptera and the genes of which have been isolated and sequenced previously : the δ -endotoxins in question are those of the strains kurstaki HD1 (19), kurstaki HD73 (20), berliner 1715 (21) and (22) Sotto (23) and aizawai IPL7 (24).

The results of these comparisons indicate that all are different in the second quarter of the molecule (amino acids 281 to 620) whereas the NH₂-terminal part (amino acids 1 to 280) and the COOH-terminal domain (amino acids 621 to 1175) of the protein are highly conserved and differ only by several amino acids. On the other hand, the δ -endotoxin corresponding to the sequence (IV) above shows considerable differences from the other δ -endotoxins both in the NH₂-terminal part (amino acids 1 to 280) and in the second quarter of the molecule (amino acids 281 to 620). The results of these comparisons indicate again that the NH₂-terminal half of the molecule (amino acids 1 to 620) which corresponds to the toxic fraction of the protein only show 46% homology with the other δ -endotoxins. The most important differences are located in the second half of the toxic part of the molecule (amino acids 281 to 620) with only 36% of identical amino acids, the NH₂-terminal part (amino acids 1 to 280) itself showing 58% of amino acids identical with the other δ -endotoxins. Such considerable differences have never been observed up to now in the NH₂-terminal part of the toxic fraction of the molecule among the δ -endotoxins active on the Lepidoptera.

In order to obtain a sequence of nucleotides entering into the framework of the invention, coding for at least the active part of a polypeptide showing a specific toxicity towards larvae of Lepidoptera of the Noctuidae family, and preferably towards S.littoralis, recourse is had, in conformity with the invention, to the following steps, namely:

- the carrying out of a molecular hybridization between, on the one hand, a nucleotide sequence of a strain of B.thuringiensis active against S.littoralis and, on the other, at least two nucleotide sequences, used as probes, derived from the 5' part of a restriction

fragment of a gene for δ -endotoxin of B.thuringiensis, this part coding for the NH₂-terminal part of the polypeptide active on the larvae of Lepidoptera, and from the 3' part of this fragment coding for the COOH part of the polypeptide,

- 5 - the isolation of the hybrid fragment,
- its cloning in a vector, followed by its purification.

In an advantageous manner, the hybridization probes utilized are obtained from a gene for the δ -endotoxin derived from the strain aizawai 7-29 coding for a protein of 130 kDa, active against P.brassicae and inactive towards S.littoralis, this gene having been
10 cloned in the recombinant plasmid pHTA2.

In an embodiment of the preceding procedure, the fragment recombined with the vector in the cloning step is elaborated from a HindIII-PstI restriction fragment derived from a single strain of B.thuringiensis, preferably aizawai 7-29. In particular, this fragment
15 is carried preferentially by the recombinant plasmid pHTA6 such as isolated with the aid of a probe constituted by a PvuII fragment of 2 kb of the plasmid pBT15-88 corresponding to the internal part of a gene for the chromosomal crystal of the strain berliner 1715,
20 starting from transforming clones containing nucleotide sequences derived from B.thuringiensis strains active against larvae of Lepidoptera, inter-alia of S.littoralis.

In another embodiment of the invention, the fragment recombined with the vector in the cloning step is elaborated from
25 several sequences of nucleotides derived from recombinant vectors containing sequences of nucleotides from at least two different strains of B.thuringiensis, possessing the same restriction maps and themselves containing all or part of the sequences of nucleotides capable of coding for a polypeptide active, in a preferential manner, against
30 S.littoralis.

In this case, the recombined fragment used in the cloning step is a fragment of about 3 kb, advantageously elaborated from a HindIII-HincII restriction fragment of about 1.1 kb derived from the entomocidus 6-01 strain and a HincII-PstI fragment of about 1.9 kb
35 from the aizawai 7-29 strain. It corresponds to a truncated gene for δ -endotoxin.

00949455-000104

The HindIII-HincII and HincII-PstI restriction fragments are carried more especially by the respective recombinant plasmids pHTE6 and pHTA6 such as isolated with the aid of the probe constituted by the PvuII fragment mentioned above.

5 The study of the toxicity towards the larvae of Lepidoptera of the bacterial strains modified with the aid of the sequences of nucleotides defined above, has made it possible to demonstrate their high toxic activity, in particular with regard to the caterpillars of S.littoralis.

10 This activity was estimated from the point of view of the specificity index corresponding to the ratio

LC50 S.littoralis

LC50 P.brassicae

15 in which "LC50" represents the lethal concentration killing 50% of the larvae in 72 hours.

The utilization of such an index makes it possible to evaluate the activity of the bacterial strains studied without having to consider the level of expression of the polypeptides.

20 The results obtained, which are reported in the examples which follow, and the values of LD50 which are deduced from them, have shown that the bacterial strains modified according to the invention show a more specific toxic activity towards S.littoralis than the native crystal proteins of the strains aizawai 7-29 or berliner 1715.

25 Therefore, the invention relates to the use of these modified strains, of recombinant vectors containing the nucleotide sequences defined above, in particular the plasmid pHT671 and the plasmid pHT71, and these sequences themselves for the elaboration of larvicidal compositions preferentially toxic towards S.littoralis.

30 The larvicidal compositions of the invention are thus characterized in that they contain an efficacious quantity of polypeptides such as defined above or expressed by the nucleotide sequences mentioned above.

35 In order to produce these polypeptides the truncated genes

00010405 "0001041

for δ -endotoxin corresponding to the nucleotide sequences of the invention are advantageously implemented.

These genes can be used to produce in excess the toxic polypeptide in microorganisms permitting the expression of the above recombinant vectors. Suitable strains of microorganisms include E.coli or also B.subtilis.

These truncated genes may be reintroduced into the strains of B.thuringiensis or into related species such as B.cereus, according to the standard techniques, for example, by transformation, conjugation or transduction. These techniques make it possible to produce the toxic polypeptide in large quantity without first having to modify the natural region of the promoter for the δ -endotoxin genes of B.thuringiensis.

This transformation may be carried out by using methods derived from the transformation of the protoplasts of B.subtilis according to (11) or of the vegetative cells of B.thuringiensis as described in (12).

The introduction of recombinant plasmids by a system of the conjugation type may be carried out by using B.thuringiensis as host strain and B.subtilis or Streptococcus faecalis as strains of the donor type by operating according to (13) and (14).

As a variant, the sequences of nucleotides are introduced into microorganisms living in the environment or in association with the plants and capable of expressing recombinant vectors containing these sequences. The introduction may be carried out in microorganisms such as Pseudomonas by working according to the procedure described in (17), by the intermediary of plasmid vectors containing the transposon Tn5 and the gene for the toxin, or Azospirillum or Rhizobium by means of the intermediary of suicide vectors derived from the plasmid RP4 and of mobilizing plasmids functional in Gram negative bacteria (for example, pRK2013) according to the procedures described in (18).

The truncated genes are alone in the strains of Bacilli or, as a variant, are associated with different δ -endotoxin genes which makes it possible to obtain crystals synthesized by these strains

specifically toxic towards given species of Noctuidae, or toxic both towards the Noctuidae and insects sensitive to other δ -endotoxins. These recombinations, carried out in vitro or in vivo with the nucleotide sequences of the invention and other δ -endotoxin genes showing different toxic specificities, lead to the construction of new genes coding for novel hybrid toxic proteins exhibiting a large spectrum of activity towards insects. These new genes and these novel proteins also enter into the framework of the invention.

In these applications, the nucleotide sequences of the invention may be transferred and expressed in plants sensitive to S.littoralis in order to diminish the devastation caused by these insects.

Among the plants to be protected, mention should be made of : cotton, clover, the tomatoe and alfalfa.

The transfer of the truncated gene into cotton plants may be carried out by transformation involving strains such as Agrobacterium as described in (15).

In addition, the invention relates to the plant cells, the plants and the seeds containing the nucleotide sequences defined above.

The plant cells according to the invention are cells, the genome of which after transformation by a non-essentially biological procedure possesses in a stable manner a sequence of nucleotides capable of expressing a polypeptide toxic towards S.littoralis, such as that defined above. The invention also relates to the plant cells derived from their division.

The plants according to the invention are plants transformed by a non-essentially biological procedure, having in particular as predator S.littoralis, the genome of which possesses in a stable manner a sequence of nucleotides such as that defined above, capable of expressing a polypeptide toxic towards S.littoralis. The plants in question are also plants derived from their reproduction, their multiplication or hybrid crosses.

In accordance with another feature, the invention relates to plants having in particular as predator S.littoralis, possessing

in addition to their initial phenotypic and genotypic characters the property of expressing a polypeptide toxic preferentially towards S.littoralis, this property resulting from the insertion in their genome by means of genetic manipulation of a sequence of nucleotides capable of expressing the said polypeptide.

In addition, the invention relates to seeds capable of giving rise to a plant such as that defined above or derived from such a plant, characterized in that they have integrated into their genome by genetic manipulation a nucleotide sequence described above.

Other characteristics and advantages of the invention will become apparent in the course of the description and in referring to the examples in which :

- figure 1 presents the restriction map of the plasmids pHTA6 and pHTE6,
- figure 2, the restriction map of a gene for a crystal protein of the aizawai 7-29 strain cloned in the plasmid pHTA2 and defining the DNA fragments which are used as probe,
- figure 3 shows the fragment of 6.6 kb cloned in pHTA6 and the result of a hybridization carried out between this fragment and the probes described in figure 2,
- figure 4, the restriction map of the plasmid pHT671, and
- figure 5, the photographs of the immunodiffusion tests.

The hybridization experiments carried out for the implementation of the invention were performed at 42°C for 24 h in a solution containing 5 x SSC, 30% formamide and 1 Denhardt (7) in the presence of the DNA probe labelled with ³²P. The filters are washed at 42°C, 20 mn, by using successively the following solutions : 5 x SSC in 30% formamide, 5 x SSC, 2 x SSC, 1 x SSC and 0.5 x SSC before drying at room temperature.

EXAMPLE 1 - Construction of a DNA sequence of about 3 kb containing a hybrid gene of an insecticidal toxin.

This construction comprises :

- 1/ the preparation of gene banks of B.thuringiensis
- 2/ the selection and characterization of transforming clones containing the genes of a crystal protein and nucleotide sequences responsible

00910405-000101

for the larvicidal activity,

3/ in vitro recombination of these sequences in a cloning vector with construction of the plasmid pHT671.

These different steps are carried out as follows :

5 1/ Preparation of gene banks of B.thuringiensis.

The total DNA of the aizawai 7-29 and entomocidus 6-01 strains of Bacillus thuringiensis is purified by using the method reported in (1) and 50 µg of each purified DNA are completely digested with the restriction enzyme PstI.

10 The DNA digested by PstI is analysed by horizontal electrophoresis on a 0.8% agarose gel and DNA fragments of a size of 5 to 8 kb are recovered from the agarose gels by electroelution in a manner described in (2).

15 The purified DNA fragments of 5-8 kb of the aizawai 7-29 strain are ligated to the DNA of the cloning vector pUC18 digested by PstI according to (3).

20 The purified DNA fragments of 5-8 kb of the entomocidus 6-01 strain are ligated to the DNA of the cloning vector pUC9 digested by PstI. The cells of E.coli JM83 are transformed with the ligation mixture as described in (4).

The transforming clones of E.coli are selected on LB medium containing 100 µg/ml of ampicillin.

2/ Isolation and characterization of the transforming clones containing the genes for a crystal protein.

25 A/ Screening of the transformed E.coli cells with the aid of an internal fragment of a gene of the crystal protein labelled with ³²P, used as probe :

30 Transforming clones containing recombinant plasmids carrying the gene for the crystal are detected by colony hybridization according to the method described in (5), by using as probe a PvuII fragment of 2 kb of the pBT 15-88 plasmid corresponding to an internal part of the gene for the crystal protein located on the chromosome of the berliner 1715 strain.

35 B/ Characterization of the recombinant plasmids present in the clones which react with the above probe.

00910405.00101

Two recombinant plasmids, pHTA6 and pHTE6, isolated respectively from gene banks constructed from the strains aizawai 7-29 and entomocidus 6-01, show a homology with this probe. In each case, a DNA fragment of about 6.6 kb was cloned.

5 The restriction map of the two plasmids is given in figure 1. The comparison of the restriction sites shows that the two DNA fragments cloned appear to be identical.

10 In order to delimit the sequences corresponding to the gene for the δ -endotoxin, different DNA fragments labelled with ^{32}P , derived from a gene of the crystal previously characterized, and cloned in the recombinant plasmid pHTA2, are utilized as probes. This latter gene for the crystal also derived from the aizawai 7-29 strain codes for a protein of 130 kd active against P.brassicae but not against S.littoralis. This type of gene possesses the same restriction map
15 as the gene for the δ -endotoxin derived from the berliner 1715 strain. In figure 2 is shown the restriction map of this gene for the crystal protein of the aizawai 7-29 strain cloned in the plasmid pHTA2. The thick lines shown above the map correspond to the fragments used as hybridization probes.

20 The plasmids pHTA6 and pHTE6 are hydrolysed by different restriction endonucleases, analysed by horizontal electrophoresis on a 0.8% agarose gel and hybridized with the different probes.

 The transfer of the DNA to nitrocellulose filters is carried out according to the method of SOUTHERN described in (6). The
25 hybridization is conducted at 42°C for 24 hours in a solution containing : 5 x SSC, 30% formamide and a 1x Denhardt mixture described in (7) in the presence of a DNA probe labelled with ^{32}P . The filters are then washed at 42°C for 20 minutes, by using successively the following solutions : 5 SSC in 50% formamide, 5 SSC, 2 SSC, 1 SSC
30 and 0.5 SSC before being dried at room temperature.

 The results of these hybridization experiments are summarized in figure 3. It appears that each extremity of the cloned DNA fragments of 6.6 kb shows a homology with the probes. The PstI-KpnI fragment of 1.5 kb reacting with the probe No. 3 corresponds to the 3' end
35 of a gene of the crystal protein present in both the aizawai 7-29

and entomocidus 6-01 strains. As pointed out in figure 3, the probes No. 1 and No. 2 corresponding to the 5' end of the gene for the δ -endotoxin of pHTA2 hybridize with the HindIII-HincII fragment of 1.1 kb contained in the plasmid pHTA6. The probe No. 3 which covers the 3' end of the gene of the δ -endotoxin of pHTA2 hybridizes with the HindIII-PstI fragment of 0.4 kb contained in the plasmid pHTA6. It should be noted that a weak hybridization signal is obtained with the probe No. 2 whereas the two other probes give easily detectable signals.

From these results, the inventors have established that the HindIII-PstI DNA fragment of 3 kb corresponds to a large part of a gene for the δ -endotoxin which commences close to the central HindIII site. It seems clear in the light of results of the hybridization experiments that the gene for the δ -endotoxin shows substantial differences from those characterized in the prior art. On the basis of these results it was decided to clone the HindIII-PstI fragment of 3 kb in the vector pUC9.

3/ Construction of the plasmid pHT 671 containing a hybrid gene of the reconstituted insecticidal toxin.

The HindIII-HincII DNA fragment of 1.1 kb derived from the plasmid pHT6 and the HincII-PstI DNA fragment of 1.9 kb derived from the plasmid pHTA6 are purified on agarose gels.

Equal amounts of the two purified DNA fragments and the DNA of pUC9 digested with HindIII and PstI are mixed and ligated. The ligation mixture is used to transform competent cells of E.coli JM83, then the transformed E.coli cells are selected on LB medium containing ampicillin. One of the interesting recombinant clones examined contains a plasmid designated by pHT671, the restriction map of which was determined and is shown in figure 4. This plasmid (pHT671) contains a DNA fragment of 3 kb inserted in the vector pUC9. This DNA sequence has the same restriction map as the HindIII-PstI fragments of 3 kb contained in the plasmids pHTA6 and pHT6, but corresponds to a reconstituted DNA molecule constructed by in vitro recombination from DNA sequences derived from the aizawai 7-29 strains on the one hand and entomocidus 6-01 on the other.

5

10

15

25

35

EXAMPLE III : Construction of a DNA sequence of about 2.7 kb containing a gene for a larvicidal toxin.

In order to achieve this construction the DNA of the aizawai 7-29 strain of B.thuringiensis was used up to the step for the production of the plasmid pHTA6 as described in Example I.

The HindIII-PstI fragment of about 2.7 kb obtained from the plasmid pHTA6 was then subcloned in the vector pUC9, previously hydrolysed by the restriction enzymes HindIII-PstI in order to give the plasmid pHT71.

EXAMPLE IV : Study of the sequence of nucleotides constituting the plasmid pHT71 coding for a polypeptide toxic towards the larvae of Lepidoptera of the family of the Noctuidae.

The HindIII-PstI fragment of 2.7 kb of pHTA6, which was subcloned in pHT71, was sequenced by means of the technique of Sanger et al. (8) using the phage M13mpl9 and the subcloning system by deletions developed by Dale et al (9). This system makes it possible to obtain M13 phages containing a series of partially overlapping DNA fragments which can be utilized for sequencing the DNA.

The sequence of nucleotides of this 2.7 kb fragment which corresponds to the chain arrangement (III) given above, was determined on the 2 DNA strands, with the exception of the last 212 nucleotides (position 2500 to 2711) which were sequenced only on a single strand.

The nucleotide sequence of this HindIII-PstI fragment has a length of 2711 nucleotides. This fragment contains the potential promoter as well as the largest part of the gene for the δ -endotoxin active on S.littoralis.

EXAMPLE V : Study of the specific toxicity of the recombinant clones of E. coli JM83 (pHT671) and JM83 (pHT71) against S.littoralis.

The toxicity of the recombinant clones of E.coli JM83 containing pHT671 and of E.coli JM83 containing pHT71 was determined by biological tests on caterpillars of the P.brassicae and S.littoralis species as described by LECADET and MARTOURET in (10). The results were compared with the specific toxicity of the native crystal proteins purified from the strains berliner 1715 and aizawai 7-29, entomocidus 6-01 B.cereus 569 (containing the plasmid pBT45, pAMB1) against the

The results obtained are reported in table 1 below.

5 In this table, for E.coli strains, the concentration 1 corresponds to a 14 hours bacterial culture concentrated 20 times, disintegrated by ultrasound ; for the B.thuringiensis strains the concentration is expressed in μg of crystal protein per μl of preparation. The toxic activity of the preparations was tested by

10 the forced ingestion with 5 μl of preparation on caterpillars at the fifth stage of development, or by a technique of free ingestion utilizing larvae at the second stage of development.

15

20

25

30

35

CONFIDENTIAL

TABLE 1

Comparative toxicity of a recombinant clone and two strains
of B. thuringiensis towards S. littoralis and P. brassicae.

5	Strains and plasmids	S.littoralis		P.brassicae		Specificity index LC50 <u>S.littoralis</u> LC50 <u>P.brassicae</u>
		LC50 2nd larval stage	LC50 5th larval stage	LC50 5th larval stage	LC50 5th larval stage	
10	JM83 (pUC18)	> 1	> 1	> 1		-
	JM83 (pHT671)	0,04	0,13	0,72		0,2
	JM83 (pHTA2)	> 1	> 1	0,03		> 30
15	JM83 (pHTA4)	> 1	> 1	> 1		-
	JM83 (pHT71)	ND	0,5	> 1		< 0,5
	berliner 1715 native crystals	ND	0,11	0,007		15,7
20	aizawai 7.29 native crystals	ND	0,02	0,04		0,5
	entomocidus 601 native crystals	ND	0,028	0,012		2,3
25	B.cereus 569 (pBT45,pAM β 1)	ND	0,38	0,054		7

3C

35

T01000 55401660

Examination of the LC50 values summarized in this table 1 shows that the protein extracts of the recombinant clones JM83 (pHT671) and JM83 (pHT71) are preferentially toxic against S.littoralis. Secondly, a comparison of the values of the specificity index shows that the larvicidal activity of the recombinant clones is more specific by a factor of 2.5 times towards S.littoralis than the native crystal proteins of the aizawai strain. Moreover, the recombinant clones of JM83 (pHT671) and JM83 (pHT71) are very active against another insect of the family of the Noctuidae, Mamestra brassicae (in the case of the clone JM83 (pHT671) for example, the LC50 value is 0.02, utilizing larvae at the second stage of development).

These two results show that the gene for the larvicidal toxin constructed and cloned in the plasmids pHT671 and pHT71 codes for a protein specifically active against S.littoralis.

Other preparations obtained from recombinant clones containing plasmids carrying genes coding for other types of δ -endotoxins (pHTA2 and pHTA4) are not active on S.littoralis : it may be seen that the plasmid pHTA2 codes for a δ -endotoxin specifically active on P.brassicae whereas the plasmid pHTA4 codes for a δ -endotoxin, the insect target for which has not yet been identified. It can also be seen that the crystalline inclusions produced by a strain of Bacillus cereus which has received the plasmid pBT45, one of the plasmids of the aizawai 7-29 strain which also carries a δ -endotoxin gene (the gene of plasmid origin of the aizawai 7-29 strain), are also specifically active on P.brassicae.

Similar results are obtained by using, in the place of crude bacterial extracts, soluble protein extracts prepared from different recombinant clones of E.coli.

On the basis of the LC50 values reported in the table above and a mean individual weight of 41 mg per L5 larva (fifth larval stage) of S.littoralis, the value of the LD50 was estimated at 2.4 μ g/gram of larva for the native crystals of the aizawai 7-29 strain.

On these same bases and on the basis of equivalence factors making it possible to pass from the total bacterial mass to the

quantity of specific proteins (about 2% of the total proteins in E.coli JM83 (pHT671), the LD50 corresponding to the toxin produced by the expression in E.coli JM83 of the gene according to the invention cloned in the plasmid pHT671, was determined and estimated at a value close to 5.5 to 6 µg/gram of larva.

10 In the two cases and particularly in the case of the ground
preparations of E.coli, the calculated values of LD50 are approximate
and higher than that of the native crystals, because it is not a
question of a purified toxin. However, these data indicate without
ambiguity that the gene expressed by pHT671 specifies a δ -endotoxin
15 exhibiting the specificity towards S.littoralis. In fact, the same
type of estimation made with extract of E.coli JM83 (pHTA2) carrying
a δ -endotoxin gene of different specificity leads to values 30 to
50 times higher than the LD50 of the soluble extracts towards
S.littoralis (135 to 350 μ g/gram of larva).

Other toxicity experiments were carried out utilizing larvae of M.brassicae, S.fruqiperda and S.littoralis at the second larval stage. The results obtained, expressed in terms of LC50 as defined for table 1, are given in table 2.

35

TABLE 2

ACTIVITY OF THE RECOMBINANT CLONES AGAINST
THE LARVAE OF INSECTS OF THE FAMILY OF THE
NOCTUIDAE: M. BRASSICAE, S. FRUGIPERDA, and
S. LITTORALIS.

STRAINS AND PLASMIDS	INSECT LARVAE AND STAGE	<u>M. BRASSICAE</u>		<u>S. FRUGIPERDA</u>		<u>S. LITTORALIS</u>	
		LC50	2nd STAGE	LC50	2nd STAGE	LC50	2nd STAGE
JM 83 (pUC10)		NT		NT		NT	
JM 83 (pHTA2)		> 1		0,51		0,9	
JM 83 (pHT671)		0,02		0,5		0,03	
JM 83 (pHT71)		ND		ND		0,03	
JM 83 (pHTA4)		> 1		0,54		> 1	

It emerges from the examination of table 2 that the crude bacterial extracts of the recombinant clone JM83 (pHT671) are toxic towards M.brassicae and S.littoralis (the values of LC50 are 0.02 and 0.03, respectively) and weakly toxic towards S.fruqiperda (LC50 of 0.5).

The extracts of the recombinant clone E.coli JM83 (pHTA2) are weakly active towards S.fruqiperda and S.littoralis and not at all toxic towards M.brassicae. The extracts of the recombinant clone JM83 (pHTA4) are not toxic towards M.brassicae and S.littoralis and are weakly toxic toward S.fruqiperda.

These results confirm the high specific toxicity of the proteins obtained from pHT71 and pHT671 towards S.littoralis and show that this class of crystal protein is also very active towards M.brassicae.

EXAMPLE VI : Study of the specificity of the polypeptides expressed by the clones formed by introduction of the plasmids pHT671 and pHT71 into E.coli.

This study was carried out owing to immuno-diffusion tests. The results are reported in figure 5 (which includes figures 5A and 5B).

The implementation of the immuno-diffusion experiment was done in conformity with the following protocol :

Soluble extracts of proteins of E.coli clones containing the plasmids pHT671, pHTA4, pHTA2 or pHT71, pUC18 were placed in the wells Nos. 2, 3, 4, 5, 6, respectively. A sample of a solubilized purified crystal of aizawai 7-29 was placed in the well No. 1 in order to serve as positive control.

In the test recorded in figure 5A an antiserum against all of the δ -endotoxins of aizawai 7-29, containing rabbit antibodies directed against the solubilized crystal proteins, was used and placed in the central well.

An immunoprecipitation line was observed in all of the cases except in the case of the extract of E.coli containing only the plasmid vector (well No. 6).

5 In the test recorded in figure 5B, the anti-serum used contained rabbit polyclonal antibodies against the crystal proteins of berliner 1715.

It may be deduced from that that the genes for the crystal isolated in pHTA4 and pHTA2 express polypeptides having antigenic determinants in common with the proteins of the crystal of berliner 1715, a strain which is not specifically active towards S.littoralis.

These results confirm those given previously with respect to the specificity of the genes isolated in the plasmids pHT71 and pHT671.

The results obtained have shown that the crystal protein represents between 7 and 10% of the total cellular proteins of E.coli JM83 (pHTA2), between 2 and 3% in E.coli JM83 (pHT671) and between 0.5 and 1% in E.coli JM83 (pHTA4) and E.coli JM83 (pHT71).

35

The literature references cited in the examples are the following :

- (1) KLIER, A.F., LECADET, M-M. and DECONDER, R., 1973, Sequential modifications of RNA polymerase during sporogenesis in Bacillus thuringiensis, Eur. J. Biochem., 36 : 317-327.
- (2) MANIATIS, T., FRITSCH, E.F., SAMBROOK, J., 1982, Molecular cloning : A laboratory manual. Cold Spring Harbor Laboratory Press, New-York
- (3) VIEIRA, J. and MESSING, J., 1982, The pUC plasmids, and M13mp7 derived system for insertion mutagenesis and sequencing with synthetic universal primers, Gene, 19 : 259-268.
- (4) LEDERBERG, E.M. and COHEN, S.N., 1974, Transformation of Salmonella thyphimurium by plasmid deoxyribonucleic acid, J. Bacteriol., 119 : 1072-1074.
- (5) GRUNSTEIN, M. and HOGNESS, D.S., 1975, Colony hybridization, a method for the isolation of cloned DNAs that contain a specific gene, Proc. Natl. Acad. Sci. U.S.A., 72 : 3961-3965.
- (6) SOUTHERN, E.M., 1975, Detection of specific sequence among DNA fragments separated by gel electrophoresis, J. Molec. Biol., 98, 503-517.
- (7) DENHARDT, D.T. 1976, A membrane filter taking for the detection of complementary DNA. Biochem. Biophys. Res. Comm., 23 : 641-646.
- (8) SANGER, F., NICKLENS, S. and COULSON, A.R., 1977, DNA sequencing with chain terminating inhibitors. Proc. Natl. Acad. Sci. U.S.A., 74 : 5463-5467.
- (9) DALE et al. (1985) A rapid single-stranded cloning strategy for producing a sequential series of overlapping clones for use in DNA, Plasmid 1985, 13 : 31-40
- (10) LECADET M.M. et MARTOURET D. 1987, Host specificity of the Bacillus thuringiensis δ -endotoxin toward

Lepidopteran species : Spodoptera littoralis Bdv and Pieris brassicae L, J. of Invert. Pathol., 49 (n° 1) : 37-48.

5 (11)CHANG et al., 1979, High frequency transformation of Bacillus subtilis protoplasts by plasmid DNA- Mol Gen Genet 168:111 115

(12)HEIERSON et al., 1987, Transformation of vegetative cells of Bacillus thuringiensis by plasmid DNA, Journal of Bacteriology, Mar.1987, p.1147-1152,

10 (13)KLIER et al., 1983, Mating between Bacillus subtilis and Bacillus thuringiensis and transfer of cloned crystal genes, Mol Gen Genet (1983) 191:257 262

(14)LERECLUS et al., 1983, Isolation of a DNA, sequence related to several plasmids from Bacillus thuringiensis after a mating involving the Streptococcus faecalis plasmid pAM β 1, Mol Gen Genet (1983) 191:307-313

15 (15)UMBECK et al., 1987, Genetically transformed cotton (Gossypium hirsutum L.) plants - Biotechnology vol.5 March 1987.

20 (16)WONG et al., 1983, transcriptional and translational start sites for the Bacillus thuringiensis crystal protein gene. J. of Biol. Chem., 258 : 1960-1967.

(17)OBUKOWICZ M. et al (1986). Tn⁵ mediated integration of the δ -endotoxin gene from B. thuringiensis into the chromosome of root colonizing Pseudomonas. J. Bacteriol., 25 168: 982-989.

(18)SIMON, R. et al, (1983). A broad host range mobilization system for in vivo genetic engineering : transposon mutagenesis in Gram-negative bacteria. Biotechnology, 1, 30 pp. 784-791.

(19)Schnepf et al, (1985) The amino acid sequence of a crystal protein from Bacillus thuringiensis deduced from the DNA base sequence. J BIOL Chem 260 : 6264-6372.

(20)Adang et al, (1985) Characterized full-length and truncated plasmid clones of the crystal protein of 35

00010405 000104

Bacillus thuringiensis subsp. kurstaki HD-73 and their toxicity to Manduca sexta. Gene 36 : 289-300.

(21)Wabiko et al,(1986) Bacillus thuringiensis entomocidal protoxin gene sequence and gene product analysis.

5 DNA 5 : 305-314.

(22) Hofte et al, (1986) Structural and functional analysis of a cloned δ -endotoxin gene of Bacillus thuringiensis berliner 1715. Eur J Biochem 161 : 273-280.

(23) Shibano et al, (1986) Complete structure of an insecticidal crystal protein gene from Bacillus thuringiensis. In : Bacillus molecular genetics and biotechnology applications. J. Ganesan, A.T., Hoch, J.A. (eds). Academic Press 307-320.

(24) Oeda et al, (1987) Nucleotide sequence of the insecticidal protein gene of Bacillus thuringiensis strain aizawai IPL7 and its high-level expression in Escherichia coli. Gene 53 : 113-119.

20

25

30

35

SECRET